

Winter Snow

Developing a Rodent Model of Adverse Menopausal Symptoms

Winter Snow¹, Lauren Federi², Stephanie Fitz³, Stephen Janasik³, Daniel Penno³, Brian Samuels², Janet Carpenter⁴, Todd Skaar⁵, Anantha Shekhar³, Philip L. Johnson³.

1. Dept. of Medicine, Ivy Tech Community College; 2. Stark Neuroscience Research Institute, 3. Dept. of Psychiatry, 4. Dept. of Adult Health, 5. Division of Clinical Pharmacology, Indiana University School of Medicine, Indianapolis, IN.

Menopause is a condition where severe depletion of estrogen levels leads to a cluster of adverse symptoms such as anxiety, cutaneous vasodilation/sudomotor "hot flashes", sleep disturbances, and appetite change (Freeman et al., 2005; Seritan et al., 2010). Previously, estrogen replacement therapy was the first line treatment for menopausal symptoms. However, it is no longer acceptable due to increased risk of cancer (Rossouw et al., 2002). Therefore there is a need for creating non-hormonal therapies to reduce the incidence of adverse menopausal-related symptoms. This is hindered by the limited understanding of menopausal symptoms and a lack of animal models of "hot flashes" (Nelson et al., 2006). Currently, the most accepted model of hot flashes is addicting female rats to morphine then inducing morphine withdrawal using naloxone (a μ -opioid receptor competitive antagonist) to provoke increases in tail temp (an indicator of cutaneous vasodilation). Yet, there is no evidence that the opioid system is disrupted in women with menopause [e.g., naloxone does not provoke "hot flashes" clinically (DeFazio et al., 1984)]. Here we induced a menopausal state by surgically removing the ovaries (OVEX) to deplete estrogen which induces a cluster of adverse menopause-like symptoms that include: 1) increased anxiety; 2) weight gain; and 3) disrupted diurnal skin and core body temperature changes. Additionally, we have developed an alternative model of "hot flashes" where administering yohimbine (an α_2 -adrenergic autoreceptor antagonist that provokes "hot flashes in menopausal women) resulted in "hot flash"-related increases in skin temp in OVEX, but not sham-OVEX, female rats.

References

- DeFazio J, Verheugen C, Chetkowski R, Nass T, Judd HL, Meldrum DR (1984) The effects of naloxone on hot flashes and gonadotropin secretion in postmenopausal women. *J Clin Endocrinol Metab* 58:578-581.
- Freeman EW, Sammel MD, Lin H, Gracia CR, Kapoor S, Ferdousi T (2005) The role of anxiety and hormonal changes in menopausal hot flashes. *Menopause* 12:258-266.
- Nelson HD, Vesco KK, Haney E, Fu R, Nedrow A, Miller J, Nicolaidis C, Walker M, Humphrey L (2006) Nonhormonal therapies for menopausal hot flashes: systematic review and meta-analysis. *JAMA* 295:2057-2071.
- Rossouw JE, Anderson GL, Prentice RL, LaCroix AZ, Kooperberg C, Stefanick ML, Jackson RD, Beresford SA, Howard BV, Johnson KC, Kotchen JM, Ockene J (2002) Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results From the Women's Health Initiative randomized controlled trial. *JAMA* 288:321-333.
- Seritan AL, Iosif AM, Park JH, Deatherage D, Sweet RL, Gold EB (2010) Self-reported anxiety,

depressive, and vasomotor symptoms: a study of perimenopausal women presenting to a specialized midlife assessment center. *Menopause* 17:410-415.